

# Opioid Program

## Pharmacist Cheat Sheet

### Identifying Patients

1. Navigate to the PrescribeCare tab and open the **Without Encounters** tab\*
2. Click on **Filters** and navigate to the **Medications** tab
3. Click on **Any Patient Score** and select **Any Opioid Risk Class**
4. Click on **Opioid Risk Class Greater Than or Equal to** and enter **5** into the text field
5. Click on **Apply**

*\*The same steps will apply for any follow-up encounters, but you will begin in the **With Encounters** tab.*

### Encounter 1: Using the Opioid Management Encounter Template

Navigate to the patient's profile and click on the **PrescribeCare** tab. Click on **Add Encounter** and select the **Opioid Management** template

*Required fields will be marked with an asterisk (\*).*

**MEDICATIONS:** Include medications pertaining to pain medication review, as well benzodiazepines and antidepressants.

**TYPE:** Select whether the encounter was an **Initial** or **Follow-Up** encounter.

### Encounter 2: Logging an MTP

Add an MTP by clicking **Edit** next to the name of the medication within the encounter. One or more of the MTPs submitted to a primary care physician (PCP) must be accepted by the PCP to be eligible for reimbursement.

<b>Pharmacist Recommendation</b>	<b>Add MTP</b>	<b>MTP Status</b>	<b>Intervention</b>	<b>Intervention Status</b>
Opioid dose lowering	Medication dose too high	Active	Recommendation to decrease medication dose	In Progress
D/C benzodiazepine	Medical contraindication	Active	Recommendation to discontinue medication	In Progress
D/C soma	Medical contraindication	Active	Recommendation to discontinue medication	In Progress

- The intervention will remain **In Progress** while waiting to hear back from the PCP
- Once a response is received from the PCP, mark the intervention status either as **Completed** if the recommendation is accepted by the PCP. Mark the intervention status as **Aborted** if the recommendation is not accepted.